## In the claims:

Cancel claims 9 and 10.

Amend claims 5 and 7 according to 37 CFR §1.121(c)(1) by replacing the existing claim with an amended claim as follows:

5. (once amended) A method according to claim 1 wherein the calcium chelating agents are selected from 4- (carboxymethyl) -2-(trimethylamino) pentane-1, 5-dicarboxylic acid, 2-(carboxymethyl) -1-(trimethylamino) butane-1, 4, dicarboxylic acid, 2-(carboxymethyl) -3-(trimethylamino) -butane-1, 4-dicarboxylic acid, ethane 1, 2-diamine N,N,N'N' tetra-acetic acid (EDTA), and codium salts of such agents and the like polydentate ligands comprising organic chelating compounds modified by addition of or substitution with a solubilizing group, e.g. a quaternary ammonium group, which is soluble in acid pH ranges, especially remaining soluble below pH4.

7. (once amended) A process according to claim 2 wherein the chelating agents include at least one of the following 4-(carboxymethyl) -2-(trimethylamino) - pentane-1, 5-dicarboxylic acid, 2-(carboxymethyl) -4-(trimethylamino) butane-1, 4, dicarboxylic acid, 2-(carboxymethyl) - 3-(trimethylamino) -butane-1, 4-dicarboxylic acid, and the like polydentate ligands comprising organic chelating compounds modified by addition of or substitution with a solubilising group, e.g. a quaternary ammonium group, which is soluble in acid pH ranges, especially remaining soluble below pH4

Add new claims 11, 12 and 13 as follows:

11. A method according to dain wherein the calcium chelating agents are selected from 4-

ay

N:\USERS\STAFF\RFP\0435\064.prl October 29, 2001 Oldfield, C. Page -4-

(carboxymethyl) -2-(trimethylamino) pentane-1, 5-dicarboxylic acid, 2-(carboxymethyl) -1- (trimethylamino) butane-1, 4, dicarboxylic acid, 2-(carboxymethyl) -3-(trimethylamino) -butane-1, 4-dicarboxylic acid, ethane 1, 2-diamine N,N,N'N' tetra-acetic acid (EDTA), and sodium salts of such agents and the like polydentate ligands comprising organic chelating compounds modified by addition of or substitution with a solubilizing group, e.g. a quaternary ammonium group, which is soluble in acid pH ranges, especially remaining soluble below pH4.

- 12. A method according to claim 11 wherein the chelating groups are selected from sulphonic and carboxylic groups.
- 13. A process according to claim 3 wherein the chelating agents include at least one of the following:4-(carboxymethyl) -2 (trimethylamino) pentane-1, 5-dicarboxylic acid, 2-(carboxymethyl) -2-(trimethylamino) butane-1, 4, dicarboxylic acid, 2-(carboxymethyl) -3-(trimethylamino) -butane-1, 4-dicarboxylic acid, and the like polydentate ligands comprising organic chelating compounds modified by addition of or substitution with a solubilising group, e.g. a quaternary ammonium group, which is soluble in acid pH ranges, especially remaining soluble below pH4.

N:\USERS\STAFF\RFP\0435\064.prl October 29, 2001